

Stored Light and Released Fiction

E.B.Alexandrov and V.S.Zapasskii

All-Russia Research Center “Vavilov State Optical Institute”, St. Petersburg, 199034 Russia

e-mail: zap@vz4943.spb.edu

It is shown that the interpretation of the experimental results reported in the publication “Storage of Light in Atomic Vapor” by D.F.Phillips et al. [1] is incorrect. The experimental observation of [1] can be consistently explained in the framework of standard concepts of the physics of optical pumping and have nothing to do with “storage of light”, or “dynamic reduction of the group velocity”, or “light pulse compression”.

I. INTRODUCTION

Our attention was attracted to paper [1] by our colleague who decided to use the results of this work to “trap, store, and release” the object beam in a telescopic system. Being aware of our experience in magneto-optics and optical pumping in alkali vapors, he asked us to comment about experimental details of this publication. So, we had to thoroughly examine this paper. The result of this examination was rather unexpected: it became evident that the authors claims on the light storage have very little to do with real physical processes in the system.

II. EXPERIMENTAL

Recall that, in the work under consideration, rubidium atoms (in zero magnetic field) were excited by a laser tuned to the transition $5^2S_{1/2}, F = 2 \rightarrow 5^2P_{1/2}, F' = 1$. The authors considered the exciting field as consisting of two coherent beams with orthogonal circular polarizations, namely, a strong σ_+ -polarized “control” beam and a weaker σ_- -polarized “signal” beam, which was turned on for a few tens of microseconds. After passing through the cell with Rb vapors, the two circularly polarized beams were detected by separate photodetectors. It is important that these two beams were treated independently, i.e., the σ_+ -component at the exit of the medium was always attributed to the control beam (σ_+ -polarized at the entrance), while the σ_- -component, to the signal beam (σ_- -polarized at the entrance). This approach contains a simple and fatal error that makes all the authors interpretation inadequate.

As experimentalists, we will analyze the observations described in [1] in general terms on the basis of simplest physical considerations to make the experimental situation as trans-

parent as possible. We will leave aside the vast theoretical background of [1] for the reason that we strongly doubt about correctness of the simplest model (the three-level Λ -scheme) used by the authors in the situation when the studied atomic line comprises 16 levels, with 11 of them being directly involved into the game through radiative and collisional processes. In addition, the strong fields used in the experiment connect 8 of these levels, thus making approximate methods inapplicable.

III. THE PICTURE OF THE OBSERVATIONS

In fact, the authors examined a response of the atomic system to a pulsed modulation of the pumping circularly polarized beam, namely, during a time interval of 10 - 30 μ s the beam was polarized elliptically. In the fundamentally nonlinear experiment (which we deal with), the medium excited by an elliptically polarized beam, with a distinguished direction in the plane of the wavefront, loses its axial symmetry (initially provided by the absence of external magnetic fields and by the axial circularly polarized pump), the circularly polarized waves cease to be eigenwaves of the medium, and the authors approach, which implies mutual independence of the circularly polarized waves in the medium, becomes invalid. Now, the “control” σ_+ -polarized beam will detect the elliptic anisotropy of the vapor. It means that the beam will exhibit elliptic birefringence and dichroism (the latter being most important), and its polarization state will vary upon its propagation through the medium. At the exit of the medium, the “control” beam will prove to be elliptically (rather than circularly) polarized and, being decomposed into the basis circular polarizations, will contribute to the signal of the second detector which is supposed to detect the “signal” σ_- -polarized beam. The same is valid for the “signal” beam, which will be detected by both detectors also. This is, briefly, what the authors observe.

IV. THREE “SURPRISING” STAGES OF THE EXPERIMENT

In more detail, the mechanism of the observed polarization dynamics looks as follows (stages 1, 2, and 3 are indicated in the figure).

Stage 1. The medium, preliminary oriented by the strong “control” beam, is subjected to a pulse of elliptically polarized excitation, which renders the medium partially aligned. In view of relatively small duration of the “signal” pulse (as compared with the spin alignment time constant), the degree of alignment continuously increases during the pulsewidth (as the integral of the pulse energy), as well as does the fraction of the “control” beam detected in the “signal” channel. This process evidently has nothing in common with the “reduction of the light group velocity” in the medium. The growth of the signal in stage 1 reflects not the shape of the leading edge of the “signal” pulse, but rather the process of accumulation

of the spin alignment. The transparency of the medium for the circularly polarized beam starts to slowly move toward its transparency for the light of elliptic polarization.

Stage 2. The light is turned off, and the alignment, thus accumulated or, more exactly, all the accumulated polarization moments remain stored in the system, relaxing with a characteristic time of about 1 ms. No energy is stored in the system and there are no grounds to be surprised that no output signal is observed as long as the light is off.

Stage 3. After the dark pause (stage 2), the “control” beam goes on detecting the residual elliptic anisotropy of the medium (partially relaxed during the pause), which is revealed, as before, in the nonorthogonality of the two circular polarizations in the medium and in branching of a fraction of the “control” beam into the “signal” channel. Now, however, the control beam, bringing strong orientation to the system, destroys the alignment much faster than it occurred in the darkness. The shape of the tail observed after the dark pause has nothing to do with that of the signal pulse. This tail reflects dynamics of the alignment destruction under the action of the control beam. This is why the duration of this tail will, in particular, depend on the control beam power, and the authors will never be able to “store” any other part of the pulse than its exponential-like tail. It may also seem curious for the authors interpretation that the total amount of the light emerging from the cell when the “stored” pulse is being “released” is smaller than when the “release” is over. This fact can be easily verified experimentally.

Thus, when we take into account the fact that the elliptically polarized light makes the medium elliptically anisotropic, all the observations became absolutely natural and trivial.

V. WHAT IS REALLY STORED

In a sense, the control beam, turned on after the pause, indeed reads out the information about the relative phase and energy of the signal beam. This information is stored in the magnitude and azimuth of the alignment and determines the phase and amplitude of the wave projected to the signal channel. But, first, the authors do not measure the phase of the signal wave and, second, what has it got to do with the storage of light? The sense of this term the authors unambiguously explain in the fragment: A pulse of light which is several kilometers in free space is compressed to a length of a few centimeters and then converted into spin excitation of a vapor of Rb atoms. After a controllable time, the process is reversed and the atomic coherence is converted back into a light pulse. In our opinion, the experiment [1] is so simple that it does leave any room for this kind of speculations. Note, in addition to all the aforesaid, that the atomic coherence carries no energy in the absence of magnetic field: the acquired spin excitation means just a sort of spatially uniform spin anisotropy of the medium. As one can easily see, in the authors terminology, any quarter-wave plate stores circularly polarized light that can be released by irradiating it with circularly polarized light

of opposite handedness. We do not find this language sensible.

VI. CONCLUSION

The interpretation of the results presented in [1] contains evident errors, and the reported experimental data (which, in themselves, seem quite correct) provide no evidence for the light “trapping, storage, and release”. In our opinion, this paper not only misleads the reader, but also arouses suspicion about other publications pertaining to the storage of light stream. This why we considered adequate evaluation of this paper important.




FIG. 1. A fragment of Fig. 2 from [1] illustrating the observed “light pulse storage” in a Rb vapor cell for a storage time of $50 \mu\text{s}$. The curves in the upper part of the figure are “calculated representations of the applied control field (dashed line) and input signal pulse (dotted line)”.

-
- [1] D.F.Phillips, A.Fleischhauer, A.Mair, R.L.Walsworth, and M.D.Lukin, Phys. Rev.Lett. 86, 783 (2001); quant-ph/0012138.